

Afric Red

Cultivar characteristics



BOTANICAL NAME	Zea Mays L.	
TYPE	Open pollinated	
AGRONOMICAL CHARACTERISTICS	Days to 50% tassel*	65 - 74
	Heat units to 50% tassel	710
	Days to physiological maturity	121 - 135
	Heat units to physiological maturity	1380
	Growth period	Medium
	Tillering**	3
	Prolificacy***	5
	Drydown period	Medium long
	Standability***	3
DISEASE TOLERANCE***	Cobrot	4
	Northern Leaf Blight	4
	Brown Rust	3
	Streak disease	5
	Grey Leaf Spot	5
REMARKS	Large cobs	

* First digit - Warm areas ** 1 - Few *** 1 - Good
 *Second digit - Cold areas ** 9 - Many *** 9 - Poor

**Variety submitted for registration (where indicated). The photo in this document only indicates the kind and not the specific variety. Technical variety information provided in this document is applicable to climatic conditions in RSA. For final cultivar recommendations in your area contact your area representative.*

For any product information outside RSA, contact the relevant KLEIN KAROO Seed Marketing (Pty) Ltd. country representative. KLEIN KAROO Seed Marketing (Pty) Ltd. supply quality seed within the tolerances laid down by the Law in South Africa. The full text of the Company's Conditions of Sale will be supplied on request.

